



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 108 363 A1**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
20.06.2001 Bulletin 2001/25

(51) Int Cl.7: **A23D 9/00**, **A23D 7/00**,
A23L 1/24

(21) Application number: **00311174.7**

(22) Date of filing: **14.12.2000**

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR
Designated Extension States:
AL LT LV MK RO SI

(72) Inventors:
• **Dartey, Clemence K.**
Ambler PA 19002-2541 (US)
• **Sox, Thomas E.**
Ambler, PA 19002 (US)

(30) Priority: **15.12.1999 US 461887**

(74) Representative: **Mercer, Christopher Paul**
Carpmaels & Ransford
43, Bloomsbury Square
London WC1A 2RA (GB)

(71) Applicant: **McNEIL-PPC, INC.**
Milltown New Jersey 08850 (US)

(54) **Long chain alcohols provided in edible oils**

(57) The present invention provides a method for incorporating long chain alcohols into edible oils. The long chain alcohol/edible oil admixture is then suitable for use

in preparation of various foods. The long chain alcohol is added by the introduction of energy and in a preferred embodiment in the absence of surfactants and emulsifiers.

BEST AVAILABLE COPY

EP 1 108 363 A1

Description**Related Applications**

5 [0001] This application is being concurrently filed with two additional applications. These related applications are entitled Encapsulated Long Chain Alcohols, European patent application No. , claiming priority from USSN 09/461 592 (Attorney's ref: P026184EP) and Long Chain Alcohols Admixed in Sterol Compounds, European patent application No. , claiming priority from USSN 09/461 607 (Attorney's ref: P026186EP).

Field of the Invention

[0002] The present invention describes a method of obtaining uniform distribution of a long chain alcohol or a mixture of such alcohols in a comestible product by dissolving or suspending the long chain alcohol in an edible oil.

Background of the Invention

[0003] Long chain alcohols are known to have beneficial effects on human health. Researchers have reported that a blend of long chain alcohols obtained from sugar cane wax was effective in lowering serum cholesterol in rats. Sho, H., Chinen, I., and Fukuda, N., J. Nutr. Sci. Vitaminol. 30:553 (1984). In addition, US Patent No.5,856,316 discloses the use of a mixture of long chain alcohols from sugar cane wax for lowering serum cholesterol in humans.

[0004] From the above disclosures it is apparent that long chain alcohols have important properties for improving the health of humans and animals. These beneficial properties include improving stamina, lowering blood cholesterol levels, and decreasing platelet aggregation. One desirable route for ingesting these alcohols is in food or beverage products. However, these long chain alcohols are very insoluble in water, and the very small amounts of these long chain alcohols needed to produce beneficial health effects makes content uniformity of long chain alcohols in a dry blending operation very difficult to achieve. Therefore, a means is needed for incorporating these long chain alcohols into food or beverage products in a manner that yields a uniform and consistent distribution of these materials in the comestible product.

[0005] Expensive and difficult measures have been disclosed to overcome these problems. For example, EP 801904 A1, describes the use of long chain alcohols (defined as alcohols having more than 20 carbon atoms) in continuous fat phase compositions containing particulate sweeteners. The incorporation of the long chain alcohols at about 0.1% to about 0.4% was reported to decrease the viscosity of confectionery coatings made with this composition. All of the described compositions contained a particulated sweetener in addition to a chocolate material and a vegetable oil.

[0006] Similarly, WO 98/47385 discloses a fat emulsion with a blend of emulsifiers created by the blending of a partial glyceride with a phospholipid and a long chain alcohol having a chain length of greater than C₂₀. The total weight of the glyceride and phospholipid in the composition must be greater than about 0.02 weight percent.

[0007] Despite the teachings of these disclosures, there is an ongoing need to easily incorporate long chain alcohols into food products so that the advantageous effects of these alcohols can be achieved.

Summary of the Invention

[0008] The present invention provides a method of providing a desirable distribution of long chain alcohols in a food or beverage product. More specifically, this process comprises dissolving or suspending the long chain alcohol in an edible oil, and then incorporating this edible oil into a food or beverage product. In one embodiment the present invention is a method for preparing a long chain alcohol in an edible oil material comprising: providing an edible oil substantially free of medium chain triglycerides composed of C₈-C₁₀ triglycerides and less than about 12 weight percent linolenic acid; providing a long chain alcohol; admixing said edible oil and long chain alcohol in the presence of an energy source such that the long chain alcohol is admixed in the oil; said long chain alcohol/edible oil admixture is stable and substantially free of an emulsifier or surfactant and having a viscosity of less than about 200 centipoise as measured at 70 °F at 60 revolutions per minute. A second embodiment of the invention provides the composition made by the method of the invention. In a third embodiment the composition of the present invention is employed to reduce the cholesterol level of a vertebrate that consumes an effective amount of the composition.

Detailed Description of the Invention

[0009] The present invention provides a method for incorporating long chain alcohols into an oil matrix without the need to add emulsifiers, surfactants or penetrant enhancers. The method includes the use of heat or mechanical energy or other suitable energy sources in order to make the long chain alcohols soluble in the lipid matrix. In a preferred

embodiment the lipid matrix is an oil, preferably an oil derived from a vegetable source. Suitable oil sources include sunflower, safflower, corn, soybean, canola, mixtures of these oils and the like.

[0010] As used herein, long chain alcohols are understood to include saturated and unsaturated alcohols which contain more than about 90 weight percent C_{20} or longer, primarily aliphatic alcohol materials. For the greatest health benefit it is preferred that the long chain alcohols be predominately, greater than 50 weight, percent octacosanol (C_{28}), preferably more than 65 percent and more preferably greater than about 70 weight percent. As used herein policosanol is understood to be a mixture of long chain alcohols ranging from C_{20} to C_{36} preferably with greater than 65 weight percent C_{28} . Common distribution and concentration ranges of the various components of policosanol are found in US Patent 5,856,316, the contents of which are incorporated by reference as set forth in its entirety. These long chain alcohols are available from various natural sources, most preferably from sugar cane wax. The long chain alcohols can also be synthesized using techniques well known in the art.

[0011] Policosanol is soluble at about 160 - 180 °F in suitable lipid matrices such as vegetable oils and fat-soluble emulsifiers, vitamins and the like. Suitable methods for incorporating policosanol or other long chain alcohol into the oil, include heating by conventional means such as: heating elements or open flame; radiation sources and other ultrasonic wave generating equipment; as well as mechanical means such as agitation, homogenization, and the like.

[0012] In a preferred embodiment of the present invention, the long chain alcohol is ground into a microcrystalline form in order to improve the stability of the oil/long chain alcohol mixture. Suitable grinding techniques include hammermill, cryogenic rotary mills and the like. The particle size of the ground material is less than about 100 microns, preferably from about 20 to about 80 and most preferably from about 30 to about 60 microns.

[0013] Typically the level of long chain alcohol incorporated into the oil is from about 0.1 to about 5 percent, preferably from about 0.5 to about 3 and most preferably from about 0.8 to about 2 weight percent. Those with skill in the art will appreciate that the concentration of the long chain alcohol will vary depending on various factors such as the dosage desired, the serving size and the solubility of the long chain alcohol in a particular oil.

[0014] One advantage of the present invention is that the long chain alcohol remains stable in the edible oil. As used herein, stable is understood to mean that the long chain alcohol does not precipitate, crystallize out, or separate when dissolved in the edible oil.

[0015] Another advantage of the present invention is that the viscosity of the policosanol-enriched oil is such that the admixture remains suitable for many applications. Unlike other disclosures, the present invention does not have to be made into a high viscosity paste before being incorporated in a comestible. The long chain alcohol/edible oil mixture has a viscosity of less than about 200 centipoise, preferably from about 10 to about 150 and most preferably from 50 to about 130 centipoise. The viscosity is measured at 70 °F using a Brookfield Viscometer Model DV-II⁺, spindle #2 at 60 revolutions per minute for 10 seconds.

[0016] Once the long chain alcohol has been suspended or dissolved into the oil matrix, those with skill in the art will readily understand how to incorporate the long chain alcohol/oil matrix into food products designed to lower cholesterol. The long chain alcohol/oil preparations can be used as a spray oil for cookies and crackers or can be used to formulate a number of cholesterol lowering food products including salad dressings, mayonnaise dressings, nutrition bars, beverages, juices, low fat ice creams, yogurts and frozen yogurts, non-dairy creamers, cheese spreads, milk products, confectioneries, chocolate-containing products such as cakes and cookies, margarine, and other spreads suitable for application on breads and the like. The long chain alcohol/oil preparations can also be formulated into pharmaceutical preparations, including tablets, soft gelatin capsules, especially those containing liquid formulations, such as suspensions, emulsions, solutions and the like.

[0017] The present invention can be provided in foods in which water is the continuous phase, such as salad dressings and mayonnaise. In a preferred embodiment of the invention the long chain alcohol/edible oil product of the present invention is incorporated into a product which has a continuous oil phase such as spreads and margarines. The long chain alcohol typically comprises from about 0.0001% to about 0.4% of the comestible product, preferably from about 0.007 to about 0.14 and most preferably from about 0.018 to about 0.071 weight percent of the comestible product.

[0018] Typically the level of the long chain alcohol is from 0.1 to about 100 milligrams/servings; preferably from about 0.5 to about 20 milligrams/serving and most preferably from about 2 to about 10 milligrams/servings.

[0019] The present invention provides an edible oil/ long chain alcohol mixture that does not require the incorporation of high levels of linolenic acid in the edible oil as was previously disclosed in the art. Further the present invention does not require the addition of triglycerides, particularly C_8 - C_{10} triglycerides; or other surfactants or emulsifiers in order to form the admixture. Because the edible oil/long chain alcohol does not need to incorporate these other agents, the admixture can be more readily formulated into other comestible products. Substantially free as used herein is understood to mean less than 1.0 weight percent, preferably less than 0.5 and most preferably less than 0.1 weight percent.

[0020] The comestible product of the present invention can be used to reduce the cholesterol level of vertebrates that consume the comestible. Vertebrates include reptiles, mammals, fish and the like, with humans being most preferred.

[0021] The following examples are provided as specific embodiments of the present invention. Other modifications

of this invention will be readily apparent to those skilled in these arts without departing from the scope of the present invention.

Example 1

[0022] One percent policosanol in a vegetable oil was prepared by adding 2.0 g policosanol (CHOLESTANOL (95 % pure sugarcane wax material from Garuda International)) to 198.0 g soybean salad oil in a 400-ml PYREX glass beaker. The beaker was placed in a waterbath on a stove. The stove was turned on and the soybean oil/policosanol mixture was stirred gently by hand with a stainless steel spatula. The policosanol/soybean oil dispersion was heated and stirred until policosanol melted and solubilized in the oil at 160 - 180 °F. The viscosity of the policosanol soybean oil preparation was measured at 170, 120, 80 and 70 °F using the Brookfield Viscometer Model DV-II+, Spindle #2 @ 60 rpm for 10 seconds. This viscosity data is presented in Table 1. Table 1 also presents results on soybean salad oil preparations containing 0, 0.1, 0.2, 0.4 and 0.6 weight % policosanol made by similar processes.

[0023] The results showed no significant differences between the viscosities of the samples at 120 °F and also at 176 °F regardless of the policosanol concentration. However, at 80 and 70 °F, the viscosities of the samples increased as the concentration of policosanol in the oil samples increased. The samples were stored at room temperature and observed the next day. The policosanol preparations developed translucent gel-like precipitates. After a few days at room temperature, the gel-like materials developed layers and gel-like precipitates. The gel-like precipitates dissolved when the samples were heated to about 180°F.

Table 1.

Effect of Policosanol Concentration on Viscosity of Soybean Oil				
% Policosanol Concentration	Viscosity @ 176 °F	Viscosity @ 120 °F	Viscosity @ 80 °F	Viscosity @ 70 °F
0	15.0 cP	26.0 cP	43.3 cP	59.5 cP
0.10	14.5 cP	27.0 cP	43.5 cP cP	64.5 cP
0.20	14.5 cP	28.0 cP	46.5 cP	82.0 cP
0.40	15.0 cP	28.0 cP	49.5 cP	119.0 cP
0.60	15.0 cP	28.0 cP	53.5 cP	112.0 cP
1.00	14.5 cP	28.0 cP	105.0 cP	129.0 cP

Example 2

[0024] A policosanol preparation from the preceding example containing 1.0 % policosanol in soybean oil was used to produce cholesterol lowering regular and light (reduced fat) margarine spreads containing plant stanol esters. The formulations are shown in Table 2. The oil phase ingredients, liquid soy bean oil (#9196 Ventura) and partially hydrogenated soybean oil (# 9494 Ventura), Canola oil, plant stanol esters, monoglycerides, lecithin and hexaglycerols were blended together in a margarine emulsion tank. The blend was mixed slowly and heated to 160 °F to melt into a clear liquid oil blend. The policosanol preparation, butter flavor, vitamins A & D blend and beta-carotene were added and blended into the oil phase.

[0025] The aqueous phase was prepared by blending potassium sorbate, ethylenediaminetetra acetic acid (EDTA), citric acid and salt in the water in an aqueous phase tank or in a stainless steel container. While stirring the oil phase at high speed, about 800 rpm, the aqueous phase was poured slowly into the oil phase to produce the margarine spread emulsion at approximately 130 ± 5 °F. The margarine was prepared by processing the emulsion through a scrape surface heat exchanger (A-unit) and then through a pin worker (B-unit) and finally through a filler unit. The finished margarine spread was filled into suitable containers. The cooling system was adjusted during the processing of the margarine spread emulsion to maintain the temperature of the finished spread at the filler unit at about 40 - 60 °F. The margarine spreads processed as indicated exhibited physical and sensory properties characteristic of regular and light margarine spreads.

Table 2.

Composition of a Typical Cholesterol-Lowering Margarine Spread Containing Policosanol.		
	Regular Margarine Spread	Light Margarine Spread
Oil Phase Ingredients	Wt. Pounds	Wt. Pounds
Liquid Soybean Oil	7.44225	23.64125
Partially Hydrogenated Soybean Oil	11.00000	0
Liquid Canola Oil	30.00000	0
Plant Stanol Esters (Raisio)	21.55000	21.55000
Mono-glycerides	0.30000	0.40000
Lecithin	0.20000	0.30000
Hexaglycerol Distearate, POLYALDO 6-2-S (Lonza)	0.10000	0.10000
Hexaglycerol Mixed Esters, CAPROL ET (A.C. Humko)	0	0.10000
Butter Flavor (Firmenich)	0.05000	0.05000
Vitamin A & D Blend (Roche)	0.00625	0.00625
Beta-Carotene	0.00150	0.00250
Policosanol/Oil Blend, (1:99 w/w) (CHOLESTANOL Policosanol from Garuda International)	4.35000	4.35000
Aqueous Phase Ingredients		
Water	22.88550	47.38300
Salt	2.00000	2.00000
Citric Acid	0.00750	0.01000
Calcium Disodium EDTA	0.00700	0.00700
Potassium Sorbate	0.10000	0.10000
TOTAL	100.00000	100.00000

[0026] Both the regular and light margarine spreads contained approximately 3.5 milligrams of policosanol per 8 gram serving size.

Example 3

[0027] The policosanol preparation, 1.0 % policosanol in soybean oil, was used to prepare ranch, French, creamy Italian and thousand island dressings containing plant stanol esters. Table 3 showed the composition of a ranch dressing produced. The process for preparing the ranch dressing in the pilot plant involved the following processing steps. The preservatives (potassium sorbate and sodium benzoate) and EDTA were dissolved in the dressing emulsion tank on a pilot plant size Charlotte Colloid Mill unit. KELTROL T xanthan gum and KELCOLOID LVF (both from Kelco) were dispersed in a portion of the soybean oil (one part gums and 2 - 5 parts oil). The gum dispersion was hydrated in the water for 10 minutes using medium speed agitation. This was followed by blending in the vinegar and lemon juice, sugar, salt, ranch seasonings, buttermilk solids and titanium dioxide. After mixing the blend at high speed for 10 minutes, the polysorbates were melted and added with the liquid egg yolk and mixed for about one minute. The policosanol oil blend was mixed into the salad oil. The stanol esters (if present) and CAPROL ET (A. C. Humko) were added to the soybean salad oil and the mixture was heated to approximately 130 °F to melt and dissolve CAPROL and stanol esters. While mixing the aqueous phase blend at high speed, the oil blend containing stanol esters (if present), CAPROL ET, policosanol oil blend, vitamin E and tocopherol preparation were added slowly to form the dressing emulsion. The coarse emulsion produced was then milled through a colloid mill with 0.02" gap opening. After the viscosity, titratable acidity, salt and pH were determined and approved, the milled dressing was pumped through a filling unit and bottled.

Table 3.

Compositions of Typical Cholesterol-Lowering Ranch Dressings Containing Policosanol with or without Stanol Esters		
Ingredient Composition	With Stanol Esters	Without Stanol Ester
	Weight (Pounds)	Weight (Pounds)
Water	33.0807	36.6479
Vinegar, 120 Grain, White distilled	7.0000	7.0000
Sugar, Fine Granulated	4.9000	4.9000
Ranch Seasoning #139 (Ventura)	4.2000	4.2000
Cultured Buttermilk Solids #64414 (Armour)	1.6000	1.6000
Salt	0.7000	0.7000
Calcium Disodium EDTA	0.0060	0.0060
Egg Yolk, Liquid, 10 % Salt	0.3600	0.3600
KELTROL T Xanthan Gum (Kelco)	0.3500	0.3500
Polysorbate 60, Tween 60	0.3000	0.3000
Lemon Juice Concentrate, 400 GPL	0.2500	0.2500
KELCOLOID LVF, (Kelco)	0.1750	0.1750
Polysorbate 80, TWEEN 80	0.1600	0.1600
Titanium Dioxide	0.1400	0.1400
Potassium Sorbate	0.0800	0.0800
Sodium Benzoate	0.0800	0.0800
Vitamin E Acetate (Roche)	0.0183	0.0183
Policosanol/soybean oil blend (1:99 w/w) (CHOLESTANOL policosanol from Garuda International)	1.2000	1.2000
DL-Alpha-Tocopherol (Roche)	0.0082	0.0082
Calcium Disodium EDTA	0.0060	0.0060
Paprika Oleoresin 1000 ASTA	0.0019	0.0019
Soybean Oil, Salad Oil	38.8186	41.1967
Stanol Esters (Raisio)	5.9453	0.0000
CAPROL ET (A. C. Humko)	0.6200	0.6200
TOTAL	100.0000	100.0000

[0028] Both dressings contained approximately, 3.5 milligram of policosanol per 30 gram serving size.

Claims

1. A process for preparing a comestible product containing high molecular weight alcohols comprising:

providing a high molecular weight alcohol;
 providing an edible oil containing less than about 12 weight percent linolenic acid that is substantially free of medium chain triglycerides composed of C₈-C₁₀ fatty acids;
 admixing said high molecular weight alcohols in said edible oil to form a high molecular weight alcohol/edible oil admixture that contains less than two weight percent high molecular weight alcohol; and combining said

admixture with other components of a comestible product.

2. The process of claim 1 wherein the high molecular weight alcohol is policosanol or octacosanol.

5 3. The process of claim 1 or claim 2 wherein the edible oil is a vegetable oil.

4. The process of any one of claims 1 to 3 wherein the comestible product is a noncontinuous oil phase product, such as a margarine, a spread, a salad dressing or a mayonnaise.

10 5. The process of any one of claims 1 to 4 wherein the amount of the long chain alcohol admixed in the edible oil comprises from 0.0001% to 0.4 weight % of the comestible product, preferably from 0.001% to 0.01% of the comestible product.

15 6. A method for preparing a long chain alcohol in an edible oil material comprising:

providing an edible oil substantially free of medium chain triglycerides composed of C₈-C₁₀ triglycerides and containing less than about 10 weight percent linolenic acid;

providing a long chain alcohol;

20 admixing said edible oil and said long chain alcohol in the presence of a energy source such that the long chain alcohol is admixed in the oil, said long chain alcohol/edible oil admixture is stable and substantially free of an emulsifier or surfactant; and having a viscosity of less than about 200 centipoise as measured at 70°F.

7. The method of claim 6 wherein the long chain alcohol and oil are heated to a temperature of from 160 to 180°F.

25 8. The method of claim 6 or claim 7 wherein the long chain alcohol is provided at a level of from 0.1 to 2 weight percent based upon the level of the oil.

9. The method of claims 6 to 8 wherein the long chain alcohol is policosanol, which preferably has an octacosanol content of greater than about 65 weight percent.

30 10. A composition comprising:

a stable edible oil/long chain alcohol admixture substantially free of an emulsifying or surfactant agent, the edible oil being substantially free of medium chain triglycerides composed of C₈-C₁₀ triglycerides and containing less than about 10 weight percent linolenic acid;

35 the admixture having a viscosity of less than about 200 centipoise as measured at 70°F.

40 11. The composition of claim 10 wherein the long chain alcohol is from 0.1 to 2 weight percent of the total composition weight.

12. The composition of claim 10 or claim 11 wherein the long chain alcohol is policosanol.

45 13. A comestible product produced by the process of any one of claims 1 to 9 or a composition of any one of claims 10 to 12 for reducing cholesterol in a vertebrate.

14. The product or composition of claim 13, which contains 0.1 to 100 milligrams of high molecular weight alcohol per serving size.

50

55



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 00 31 1174

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
Y	DATABASE WPI Section Ch, Week 198618 Derwent Publications Ltd., London, GB; Class D13, AN 1986-117166 XP002042067 -& JP 61 058536 A (NIPPON OILS & FATS CO LTD), 25 March 1986 (1986-03-25) * abstract *	1-14	A23D9/00 A23D7/00 A23L1/24
Y	PATENT ABSTRACTS OF JAPAN vol. 009, no. 180 (C-293), 25 July 1985 (1985-07-25) -& JP 60 049752 A (SANYUU SHOJI KK), 19 March 1985 (1985-03-19) * abstract *	1-14	
D,A	EP 0 901 804 A (LODERS CROKLAAN BV) 17 March 1999 (1999-03-17) * page 2, line 26 - line 29 * * claims 1-3 *	1-3,5, 10,13,14	
D,A	WO 98 47385 A (HAILES ANNE ;UNILEVER PLC (GB); VERMAAS LEO FRANS (NL); ZWEMMER OL) 29 October 1998 (1998-10-29) * examples I-III * * claims 1,4 *	1-5,10	TECHNICAL FIELDS SEARCHED (Int.Cl.7) A23D A23L
A	DATABASE WPI Section Ch, Week 199610 Derwent Publications Ltd., London, GB; Class D13, AN 1996-095736 XP002163880 & KR 9 404 531 B (OTTOGI FOODS CO LTD), 25 May 1994 (1994-05-25) * abstract *	1,6,10	
-/--			
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 26 March 2001	Examiner Dekeirel, M
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application I : document cited for other reasons & : member of the same patent family, corresponding document	

PROTON 1500 03 02 (P4001)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 00 31 1174

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
A	PATENT ABSTRACTS OF JAPAN vol. 012, no. 092 (C-483), 25 March 1988 (1988-03-25) -& JP 62 224258 A (JUN KAWAI), 2 October 1987 (1987-10-02) * abstract *	10	
A	PATENT ABSTRACTS OF JAPAN vol. 018, no. 316 (C-1213), 16 June 1994 (1994-06-16) -& JP 06 070717 A (NIPPON OIL & FATS CO LTD), 15 March 1994 (1994-03-15) * abstract *	10	
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 26 March 2001	Examiner Dekeirel, M
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 1503 03 82 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 00 31 1174

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

26-03-2001

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
JP 61058536	A	25-03-1986	NONE	
JP 60049752	A	19-03-1985	NONE	
EP 0901804	A	17-03-1999	NONE	
WO 9847385	A	29-10-1998	AU 7430698 A EP 0977489 A	13-11-1998 09-02-2000
KR 9404531	B	25-05-1994	NONE	
JP 62224258	A	02-10-1987	NONE	
JP 06070717	A	15-03-1994	NONE	

EP FORM P4468

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☒ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.